

Stretch Mark Therapy No. 2

(i.e. pregnancy)

1- 2 times a week for 4 - 6 weeks

For better results the two Stretch Mark therapies can alternate.

Therapy Procedure

<p style="text-align: center;">Brushes</p> <p>Cleansing Oil - H0501040</p>	<p>Use the brushes (hard brush) in circular clockwise movements, until the skin reddens lightly.</p> <p>Continuing apply Cleansing Oil</p>
<p style="text-align: center;">Thermo Mask</p> <p>Thermo Packung - H0200134 Actif Gel - H0208882</p>	<p>Apply Thermo Packung (H0200134) in the treatment area but leave it on only for 3 minutes. Remove the mask and cover with towels.</p> <div style="border: 1px solid black; padding: 5px; margin: 10px 0;"> <p>Attention: Avoid the armpits area and the area on and around the nipples.</p> </div> <p>For Sensitive Skin: Instead of the Thermo Packung use the Actif Gel</p> <p>During the treatment the area where we apply the Thermo Packung reddens. This is part of the therapy.</p>
<p style="text-align: center;">Multipulle Colastin Serum</p>	<p>Apply on the treatment area the Multipulle Colastin Serum 1-2 ml and massage lightly.</p>
<p style="text-align: center;">Iontocutin N Cream</p>	<p>Apply Iontocutin N Cream and massage lightly for about 20 minutes.</p>
<p style="text-align: center;">Bandages</p> <p>Bandages - H0208809</p>	<p>Cover the treated area with warm bandages and cover with a dry towel. Let it take effect for about 20 minutes</p>
<p style="text-align: center;">Total Therapy Time</p>	<p>About 45 minutes</p>
<p style="text-align: center;">Essential For Home →</p> <p>Collastin ampoules- H0303317 Phyto Collastin Solution 30 ml H0100155 Iontocutin N Cream</p>	<p>At home the client must use Collastin ampoules followed by Iontocutin Cream for 10 days. In addition the client can also use the Phyto Collastin Solution 30 ml for even better results.</p>